

## National Childhood Immunisation Schedule, Singapore

Vaccination against	Birth	1 Month	3 months	4 months	5 months	6 months	12 months	15 months	18 months	6-7 years^	10-11 years^^
Tuberculosis	BCG										
Hepatitis B	HepB (D1)	HepB (D2)				HepB (D3)†					
Diphtheria, Tetanus, Pertussis			DTaP (D1)	DTaP (D2)	DTaP (D3)				DTaP (B1)		Tdap (B2)
Poliovirus			IPV (D1)	IPV (D2)	IPV (D3)				IPV (B1)		OPV (B2)
<i>Haemophilus influenzae type b</i>			Hib (D1)	Hib (D2)	Hib (D3)				Hib (B1)		
Measles, Mumps, Rubella							MMR (D1)		MMR (D2)##		
Pneumococcal Disease			PCV (D1)		PCV (D2)		PCV (B1)				
Human Papillomavirus		<b>Recommended for <u>females 9 to 26 years</u>; three doses are required at intervals of 0, 2, 6 months</b>									

**Notes:**

- BCG Bacillus Calmette-Guérin vaccine
  - HepB Hepatitis B vaccine
  - Hib *Haemophilus influenzae* type b vaccine
  - DTaP Paediatric diphtheria and tetanus toxoid and acellular pertussis vaccine
  - Tdap Tetanus toxoid, reduced diphtheria toxoid and acellular pertussis vaccine
  - MMR Measles, mumps, and rubella vaccine
  - OPV Oral polio vaccine
  - IPV Inactivated polio vaccine
  - PCV Pneumococcal conjugate vaccine
  - D1/D2/D3 1<sup>st</sup> dose, 2<sup>nd</sup> dose, 3<sup>rd</sup> dose
  - B1/B2/B3 1<sup>st</sup> booster, 2<sup>nd</sup> booster, 3<sup>rd</sup> booster
  - ^ Primary 1
  - ^^ Primary 5
- # 3<sup>rd</sup> dose of HepB can be given at the same time as the 3<sup>rd</sup> dose of DTaP, IPV, and Hib for the convenience of parents.
  - ## 2<sup>nd</sup> dose of MMR can be given between 15-18 months